

Miami, Florida – 1999 (N=208)

Figure A. Age of GISP participants, in years, 1999




Figure B. Race/ethnicity of GISP participants, 1999




Figure C. Percentage of GISP participants identifying as men who have sex with men, 1988–1999






Figure D. Reason for visit among GISP participants, 1999



Miami, Florida – 1999 (N=208)

Figure E. Previous episode of gonorrhea among GISP participants, 1991 – 1999



*Data first collected in 1991.

**Data first collected in 1992.

Note: Data points not shown when >30% data missing.

Figure F. Drugs used to treat gonorrhea among GISP participants, 1999




Figure G. Drugs used to treat chlamydia trachomatis infection among GISP participants, 1999





Figure H. Resistance to penicillin and tetracycline among GISP isolates, 1999



Miami, Florida – 1999 (N = 208)

Figure I. Decreased susceptibility to ceftriaxone among GISP isolates, 1988 – 1999

No isolates with decreased susceptibility to ceftriaxone have been identified at this clinic.

Figure J. Decreased susceptibility to cefixime among GISP isolates, 1992 – 1999

No isolates with decreased susceptibility to cefixime have been identified at this clinic.

Figure K. Intermediate resistance and resistance to ciprofloxacin among GISP isolates, 1990 – 1999




Figure L. Decreased susceptibility to azithromycin among GISP isolates, 1992 – 1999

No isolates with decreased susceptibility to azithromycin have been identified at this clinic.

Note: Susceptibility to ciprofloxacin first measured in 1990.